

## Contents

1. Product description
2. Background information and applications
3. Protocols
  - 3.1 Preparations
  - 3.2 Culture conditions
  - 3.3 Instructions for use
    - 3.3.1 Thawing of cells
    - 3.3.2 Routine cultivation and cell expansion
    - 3.3.3 Stepwise adaption from serum-containing cultures
    - 3.3.4 Bioreactor cultivation
    - 3.3.5 Freezing of cells
4. Ordering information

## 1. Product description

### Components and specifications

HEK GM medium  
(for preparation of liquid media from powder  
see Solubilization Protocol)

without L-glutamine  
with growth hormone

Chemically defined  
Free of animal-derived components  
Free of proteins

### Storage

Store protected from light at 2–8 °C. Do not freeze.

### Intended use

Intended for *in vitro* research and manufacturing processes **only**. Do not use for injection or infusion!

## 2. Background information and applications

HEK GM is a complete chemically-defined, animal-component-free and protein-free medium. HEK GM was developed by Xell for high-performance cultivation of HEK and other human cell lines. HEK GM supports cell growth and production of recombinant proteins and antibodies in suspension culture. It can be used in research or in manufacturing applications.

## 3. Protocols

### 3.1 Preparations

All procedures should be carried out using sterile techniques in a biosafety cabinet.

HEK GM is formulated without L-glutamine. For applications requiring this amino acid, supplement with 6-8 mM L-glutamine prior to use. Supplementation of L-glutamine directly to the culture is recommended.

*Note: No supplementation with e.g. Pluronic® F68 is necessary to maintain cells in suspension.*

### 3.2 Culture conditions

Cultures should be maintained at 37 °C. For cultivation in an incubator, a 5% CO<sub>2</sub> atmosphere is necessary.

Parameter	Value[-]
Shaker diameter	5 cm
Shaker speed	125-185 rpm
Temperature	37°C
CO <sub>2</sub>	5%

**Table 1:** Recommended culture conditions for use of Xell media and feed products.

Using the set-up listed in table 1, the working volume of different polycarbonate Erlenmeyer shake flask sizes was determined (table 2). For cell lines with strong aggregation, baffled shake flasks may be used. For this setup, a reduction of the shaking speed might be necessary.

Size of shaker [mL]	Shape [-]	Working volume [mL]
125	plain, vent cap	20 - 50
250	plain, vent cap	80 - 150
500	plain, vent cap	200 - 300
1000	plain, vent cap	400 - 600

**Table 2:** Recommended culture working volumes for use of Xell media and feed products in various shake flask sizes.

### 3.3 Instructions for use

#### 3.3.1 Thawing of cells

- 1) Quickly thaw a vial of frozen cells in a 37 °C water bath.
- 2) Transfer the cells aseptically to a centrifugation tube containing 10 mL of HEK GM.
- 3) Centrifuge cell suspension at 115×g for 5 minutes.
- 4) Aspirate supernatant completely and discard.
- 5) Resuspend the cells in 10 mL HEK GM per vial.
- 6) Adjust viable cell density to 5-10×10<sup>5</sup> cells/mL by medium addition and transfer cell suspension into an

agitated or stationary cultivation system (e.g. T-75 tissue culture flask, 125 mL polycarbonate Erlenmeyer flask, or 50 mL filter tube).

- 7) Count the cells after 24-48 hours for assessment of cell density and viability.
- 8) Adjust cell density to  $3-6 \times 10^5$  cells/mL. \*
- 9) Proceed with routine cultivation.

\* Depending on the cell line, the target inoculation cell density can be lower.

### 3.3.2 Routine cultivation and cell expansion

- 1) Pre-equilibrate a sufficient amount of medium in a polycarbonate Erlenmeyer shake flask (Parameters listed in tables 1 and 2) for 1 hour. \*\*
- 2) Determine viable cell density in the pre-culture.
- 3) Depending on the inoculation volume, remove medium from the shake flask to reach the target working volume after inoculation. Final working volume of given shaker size is listed in table 2.
- 4) Seed cells at a target inoculation cell density of  $3 \times 10^5$  cells/mL (operational range  $2-5 \times 10^5$  cells/mL).
- 5) Incubate the culture according to the conditions listed in table 1.
- 6) Routinely passage the culture when viable cell densities between  $15-40 \times 10^5$  cells/mL are reached. Typical duration time for the culture is 3-4 days.
- 7) If cell density is too low or cells do not grow in adaption phase, centrifuge the culture and exchange the medium without dilution after 4 days.

\*\* Depending on cell line, the pre-equilibration of medium might be not necessary. For some cell lines the use of 2-8°C cold culture medium directly from refrigerator was found to be beneficial. This procedure eliminates handling variations of the medium in the pre-equilibration phase of the medium.

### 3.3.3 Stepwise adaptation from serum-containing cultures

- 1) Expand the culture in serum-containing standard medium.
- 2) Centrifuge a sufficient number of cells for inoculation of suspension culture with  $4-6 \times 10^5$  cells/mL at  $115 \times g$  for 5 minutes.
- 3) Resuspend cells in Xell medium (if necessary, include 6-8 mM L-glutamine) and 2 % fetal bovine serum (FBS).
- 4) Passage cells or change medium by centrifugation every two to four days depending on cell density.
- 5) Reduce serum concentration to 0.5 % after at least three passages.
- 6) Passage cells or change media by centrifugation every two to four days depending on cell density.
- 7) Reduce serum concentration to 0 % after two to four passages.
- 8) Continue cultures until viabilities stabilize at > 90 %.
- 9) Adapted cells should be inoculated at  $2-5 \times 10^5$  cells/mL in Xell medium for optimal performance. Cultures should be diluted every three or four days. Due to aggregation of HEK cells, cultures should be stirred or shaken, using spinner bottles, shaker flasks or similar cultivation systems.

### 3.3.4 Bioreactor cultivation

For best performance the inoculation density in bioreactor should be in the range of  $4-6 \times 10^5$  cells/mL in Xell medium. Suggested starting parameters for bioreactor cultivations of HEK cells using Xell medium are pH 7.0-7.5, 40% DO, and a temperature of 37 °C.

*Note: No supplementation with e.g. Pluronic® F68 is necessary to maintain cells in suspension.*

### 3.3.5 Freezing of cells

Cells can be frozen in HEK GM medium without the use of serum.

- 1) Choose a well-growing culture with viabilities above 90 %.
- 2) Prepare a freezing medium consisting of 90 % HEK GM medium and 10 % dimethyl sulfoxide (DMSO; cell culture grade).
- 3) Cool down the freezing medium to 2-8 °C.
- 4) Centrifuge the cells at  $115 \times g$  for 5 minutes.
- 5) Aspirate supernatant completely.
- 6) Resuspend the cells in freezing medium at  $1 \times 10^7$  cells/mL.
- 7) Rapidly transfer 1.5 mL of this suspension to sterile cryovials.
- 8) Place the vials in a pre-cooled (2-8 °C) freezing module and store the modules including the vials for 24 hours at -80 °C.
- 9) Transfer the cryovials to a -140 °C to -196 °C system for long time storage.

## 4. Ordering information

Product	Application	Order No.
HEK GM	base medium and protein production for various human cell lines, e.g. HEK cells	851
HEK TF	transfection medium, suitable for growth, transfection/infection and production for various HEK cells and other human cell lines; esp. for transfection	861
HEK ViP NB	Transfection and production medium for HEK cells with a <b>basic nutrient</b> level	891
HEK ViP NX	Transfection and production medium for HEK cells with a <b>high nutrient</b> level	892
Basic Feed	feed supplement for CHO, HEK, hybridoma, et al.	1092
HEK FS	Feed supplement for various human cell lines, e.g. HEK	871

**Table 3:** HEK products by Xell

Place orders: [order@xell.de](mailto:order@xell.de)

For further information or assistance contact us.

[www.xell.de](http://www.xell.de)  
[info@xell.de](mailto:info@xell.de)

Xell AG  
Alte Verler Str. 1  
33689 Bielefeld  
Germany

Fon: +49 (0)521 96989-200  
Fax: +49 (0)521 96989-201

